CUMULATIVE INDEX 1994

Volume 10

January
April
July
PREDICTING INTENSIVE CARE UNIT OUTCOME, pages 1–246
IMAGING IN THE INTENSIVE CARE UNIT, pages 247–459
July
TRAUMA AND PRE-EXISTING DISEASE: PART II SPECIFIC
MANAGEMENT CONCERNS, pages 461–649
October
IMPROVING PATIENT TOLERANCE OF MECHANICAL
VENTILATION, pages 651–883

Note: Page numbers of issue and article titles are in boldface type.

Abdomen, aortic aneurysm of, ultrasonography of, 354–357 CT of, 321–339. See also specific diseases. infection of, scintigraphy for, 387 plain film radiographic examination of, 277–296

ultrasonography of, **297–319**. See also specific sites.

Abdominal aortic rupture, CT of, 335 Abruptio placentae, in pregnant trauma

patients, 630 Abscess(es), appendiceal, CT of, 333–334 colonic diverticular, CT of, 334

extraperitoneal, CT of, 332–334

hepatic, CT of, 322–323 ultrasonographic detection of, 304–305

intra-abdominal, ultrasonographic detection of, 315–316

intraperitoneal, CT of, 332-334 mediastinal, chest CT for, 271

pelvic, MRI of, 409 percutaneous management of, 445–446 renal, CT of, 329–330

splenic, CT of, 328–329

Abscess cavity, plain film radiographic examination of, 285, 288, 289

Acetylcholine (ACH), in neural communication, 781

Acidosis, lactic, in shock, 181–183
Acquired immunodeficiency syndrome

Acquired immunodeficiency syndrome (AIDS), MRI of, 413–414

Actigraphy, in sleep measurement, 684 Acute myopathy syndrome, 805–806 Acute Physiology and Chronic Health Evaluation. See APACHE. Addison's disease, described, 543–545

Adjustment disorder with anxiety, in ICU, 677

Adrenal insufficiency, described, 543–545 Adult respiratory distress syndrome (ARDS), acute, mortality following, 206

outcome following, predictors of, 207–208

outcome following, 206–212 functional, 208–210

radiographic evaluation of, 261-263

Age, as factor in peripheral arterial disease, 567

as factor in trauma patients with pre-existing cardiac disease, 461–463 Agitation, defined, 661

drugs for, 709-731

life-threatening episodes of, 750–753

AIDS. See Acquired immunodeficiency syndrome.

Airway obstruction, in near-death experience, 655–656

in trauma patients with chronic pulmonary disease, 514-515

Alcohol, use of, classification of, 605 Alcohol abuse, in brain-injured trauma

patients, 636 motor vehicle accidents and, 595, 596

in trauma patients, 595-611 acute intoxication, 596-597 Alcohol abuse (Continued) chronic abuse, and withdrawal syndrome, 597–598

Alcohol withdrawal, delirium tremens with, 678, 751

Alpha₂-adrenoreceptors, for hyperadrenergic hyperfunction, 743 Amniotic fluid embolus, in pregnant

trauma patients, 630

Amrinone, in trauma patients with preexisting cardiac disease, 491

Analgesia, patient-controlled, for pain in mechanically ventilated patients, 769

Analgesics, in organ system failure, 745–746

Anesthesia/anesthetics, in the ICU, 437–438

local, with opioids, combined infusions of, for pain in mechanically ventilated patients, 773

for pain in mechanically ventilated patients, 770

in trauma patients with pre-existing cardiac disease, 484, 488–489

Aneurysm, aortic, abdominal, ultrasonography of, 354–357 mycotic, of the ascending aorta, 404–405

Angina, defined, 475 evaluation and management, 475

pre-existing, in trauma patients, 466 Anti-inflammatory drugs, nonsteroidal (NSAIDs), for pain in mechanically ventilated patients, 774

Antibiotic(s), in trauma patients with preexisting peripheral arterial disease, 578–579

Anxiety, adjustment disorder with, in ICU, 677

causes, 675 defined, 661, 674 in ICU, 673-675

weaning and, 678-679

Aorta, ascending, aneurysm of, 404–405 injury to, in trauma patients with pre-existing peripheral arterial disease, reconstruction for, 584–588

Aortic aneurysm, abdominal, ultrasonography of, 354–357

Aortic dissection, CT for, 271

Aortic insufficiency, in trauma patients
with pre-existing cardiac disease,
treatment, 493

Aortic stenosis, in trauma patients with pre-existing cardiac disease, treatment, 492

APACHE, 59–60 ethical appropriateness of, 233–236 limitations of, 235–236 in prediction of ICU outcome, case against using, 117–125, 127–128, 129–134

properties of, 3-4 uses of, 234-235

APACHE II, errors and bias in, 60-64 APACHE III, 94-98

admission in, precipitating reasons for,

in the Cleveland project, 151–153 components of, 94, 95

errors and bias in, 64-68

score ranges of, 94

Appendicitis, CT of, 333-334

Appendicoliths, plain film radiographic examination of, 293, 294

Apprehension, defined, 661

ARDS. See Adult respiratory distress syndrome.

Arousal, autonomic, defined, 661 Arrhythmia(s), pre-existing, in trauma patients, 463–464

in trauma patients, 469-470 treatment, 493-498

Artery(ies), axillary, injury to, in trauma patients with pre-existing peripheral arterial disease, treatment, 584

brachiocephalic, injury to, in trauma patients with pre-existing peripheral arterial disease, treatment, 583

extrathoracic subclavian, in trauma patients with pre-existing peripheral arterial disease, treatment, 583–584

injury to, in trauma patients with pre-existing peripheral arterial disease, treatment, 582–583

peripheral, disease of. See Peripheral arterial disease (PAD).

pulmonary, catheters in, placement and misplacement of, 427–430

Ascites, in patients with pre-existing liver disease, 560–563

Aspiration, radiographic evaluation of, 260 ultrasound-guided, 315–316

Atelectasis, radiographic evaluation of, 256-257

Atracurium, description and uses of, 787, 790

for respiratory failure, 835, 836 indications, 840

Atrial fibrillation, in trauma patients, treatment, 494–495

Atrial flutter, in trauma patients, treatment, 494-495

Autonomic arousal, defined, 661

Axillary artery, injury to, in trauma patients with pre-existing peripheral arterial disease, treatment, 584 Barbiturate(s), abuse of, in trauma patients, 601 clinical indications, 717

pharmacokinetics of, 716 pharmacology of 716–717

pharmacology of, 716–717 physiology of, 716 side effects, 716

Benzodiazepines, abuse of, in trauma patients, 601–602

clinical indications, 714–715 for delirium tremens, 753

for hyperadrenergic hyperfunction, 738–743

pharmacokinetics of, 713 pharmacology of, 711–715

receptor physiology of, 711–712 side effects, 713–714

for sleep disturbances in ICU, 690-691 Bifascicular block, in trauma patients, for

atrial fibrillation/flutter, 498
Biliary tree, calcifications in, plain film
radiographic examination of, 294

gas in, plain film radiographic examination of, 288–289, 291

ultrasonography of, 297–303 Biofeedback, for patient distress during

mechanical ventilation, 103
Biopsy, muscle, in evaluation of muscle

weakness following neuromuscular blockade, 804

nerve, in evaluation of muscle weakness following neuromuscular blockade, 804

Bleeding, gastrointestinal, radionuclide evaluation of, 375–378

in trauma patients with pre-existing peripheral arterial disease, treatment, 579–580

Blood gases, arterial, for trauma patients with chronic pulmonary disease, 509-510

Blood pressure, intra-arterial, in trauma patients with pre-existing cardiac disease, 481

Blood transfusion, in traumatized renal failure patients, 528–529

Blood volume, systemic, evaluation of, chest radiography for, 251

Bowel, obstruction of, plain film radiographic examination of, 279–292

Bowel gas patterns, plain film rac...graphic examination of, 279–292

Brachial plexus blocks, "single-shot," 775 Brachiocephalic artery(ies), injury to, in trauma patients with pre-existing peripheral arterial disease, treatment, 583

Bradyarrhythmia(s), in trauma patients, treatment, 497

Bradycardia, junctional, in trauma patients, treatment, 497

sinus, in trauma patients, treatment, 497 Brain, scintigraphy of, 378–380

Brain death, radionuclide scanning in, 379
Brain failure, cerebral function monitoring
for, 756–759

dopaminergic imbalances in, 737–738 in ICU, 734–735

neurobiology of, 733-734

noradrenergic hyperfunction in, treatment, 738–743

symptoms, 735-736

treatment, cholecystokinin antagonists, 760–761

drugs in, 733–765. See also Organ system failure.

serotonin in, 759-760

Brain injury, irreversible, prediction of, future directions in, 225-226

traumatic, in patients with neurologic and psychiatric disorders, risk of, 635–638

Breathing, long-term ventilation effects on, 852–856

Brief-MAST, 606-607

Bupivacaine, for pain in mechanically ventilated patients, 770–771

Buspirone, in brain failure and delirium, 760

Cacification(s), abdominal, plain film radiographic examination of, 293–294 CAGE Questionnaire, 606

Calf, veins of, vascular ultrasonographic evaluation of, 351–353

Calibration, defined, 56–57 in probability model, 101–102

Candidiasis, hepatic, ultrasonographic detection of, 305

Carbon dioxide, end-tidal, in cardiac arrest, 189–191

Carbon monoxide poisoning, in pregnant trauma patients, 631

Cardiac arrest, coronary perfusion pressure in, 188–189

end-tidal carbon dioxide in, 189–191 ICU outcome related to, 188–191

Cardiac disease. See Heart disease.

Cardiac function, in trauma patients with pre-existing peripheral arterial disease, 574-575

Cardiac output, in trauma patients with pre-existing cardiac disease, 482

Cardiopulmonary resuscitation (CPR), coma following, prognosis, 221–222 Cardiovascular system, abnormalities of, chest CT for, 271-274 in patients with pre-existing liver disease, 559-560

benzodiazepines effects on, 714 evaluation of, chest radiography for, 250-255

narcotic effects on, 724

in obese trauma patients, 616-617 pregnancy-induced changes in, 624

Carotid artery(ies), injury to, in trauma patients with pre-existing peripheral arterial disease, treatment, 582-583

Catecholamine(s), in trauma patients with pre-existing cardiac disease, 491

Catheter(s), central venous, insertion of, 438-441

placement and misplacement of, 424-427

epidural, in mechanically ventilated patients, in pain management, 770-773 fragments of, retrieval of, in the ICU,

441-443 placement and misplacement of, 417-436 See also specific types of catheters.

pulmonary artery, placement and mis-placement of, 427-430 Catheterization, arterial, complications of,

ultrasonography of, 357-359 Cecum, distention of, plain film radiographic examination of, 281

Central nervous system (CNS), infections of, MRI of, 413-414

Central nervous system (CNS) function, in trauma patients with pre-existing

peripheral arterial disease, 575 Central nervous system (CNS) injury. See also specific types of injury.

future directions in, 225-226 ICU outcome following, 217-228

Central venous pressure, in trauma patients with pre-existing cardiac disease, 481

Cerebral function monitor, in sleep measurement, 684

Cerebral function monitoring, in brain failure and delirium, 756-759

Cerebrovascular disease, in trauma patients, 569-570

Cesarean section, perimortem, in pregnancy trauma patients, 631

Chest, CT of, in the ICU, 267-275 Chest radiography, in ARDS evaluation,

261-263 in aspiration evaluation, 260 devices on, 249-250

in evaluation of cardiovascular status, 250-255

in evaluation of pulmonary disease, 256-260

film analysis, 248

portable, uses of, 247-248

Chest trauma, CT for, 271

Cholangitis, ascending, CT of, 322

Cholecystitis, acalculus, ultrasonographic detection of, 301

acute, CT of, 321-322 scintigraphy in, 371-373

acute calculus, ultrasonographic detection of, 298-301

emphysematous, plain film radiographic examination of, 289

gangrenous, ultrasonographic detection of, 301

Cholecystokinin antagonists, in brain failure and delirium, 760-761

Cholecystokinin receptors, in brain failure and delirium, 761

Choledocholithiasis, ultrasonographic detection of, 302-303

Chronic obstructive pulmonary disease (COPD), acute respiratory failure in, outcome following, 197-215

postoperative complications related to, 512-513

Cirrhosis, ascites in, 560-563 in trauma patients, 558

Cleveland, reimbursement strategies in, severity scoring systems for, 148-150 APACHE III in, 151-153

paying for, 154-155 Clinical abstractions, 48

Clonidine, for hyperadrenergic hyperfunction, 743

for life-threatening delirium, 754

CMV. See Cytomegalovirus.

CNS. See Central nervous system. Coagulation, diabetes mellitus and, 542-543

Coagulopathy, in patients with pre-existing liver disease, 560

Cocaine, abuse of, in trauma patients, 602-604

Cognitive-behavioral strategies, for patient distress during mechanical ventilation, 700-704

Cognitive distortion, defined, 661

Colitis, plain film radiographic examination of, 282-283

Colon, obstruction of, plain film radiographic examination of, 280

Coma, defined, 661

following cardiopulmonary resuscitation, prognosis, 221-222

following drug overdose, prognosis, 221

following stroke, prognosis, 219-221 nontraumatic, prognosis, 218-222

poor outcome following, predictors of, 225

traumatic, prognosis, 222-224

Communication, with intubated patients, 696–698

Computed tomography (CT), abdominal, 321–339. See also specific diseases. for abscesses, 332–334

in acute hepatobiliary inflammatory disease, 321–323

in acute pancreatic inflammatory disease, 324–328

in intestinal diseases, 334-335 for jaundice, 323-324

in kidney disease, 329-332 for spleen, 328-329

chest, for cardiovascular abnormalities, 271–274

in the ICU, 267-275 in lung disease, 268-269

for mediastinal abnormalities, 271 in pleural disease, 269–270

Conduction defects, in trauma patients, treatment, 497–498

Control, loss of, in near-death experience, 652–653

Coronary artery disease, in trauma patients, 569

Coronary care unit (CCU), use of severity of disease systems in, 175

Coronary perfusion pressure, in cardiac arrest, 188–189

Corticosteroid(s), for trauma patients with chronic pulmonary disease, 515

Cost-containment, severity models for, 103 Critical care, history of, 157–159 Critical care medicine, effects on ICU

outcome, 165–168 Critical care research, clinical databases

and, 46–49

Cross-clamping aortic in trauma patien

Cross-clamping, aortic, in trauma patients with pre-existing cardiac disease, 484

CT. See Computed tomography. Curare, description of, 790

for respiratory failure, 835, 836 uses of, 790

Cystitis, emphysematous, plain film radiographic examination of, 290, 292 Cytokine markers, in shock, 184–185

Cytomegalovirus (CMV) pneumonia, 257

Data, acquisition of, 43–45 recommendations for, 49 definition and validation of, 42–43 recommendations for, 49 management, 45–46

recommendations for, 50

manipulation of, 45-46

objective quality assurance, 157-177 need for, 157-159

storage of, recommendations for, 49-50

Data dictionary, 43 Data glut, 47

Data model, 48

Data poverty, 47

Database(s), clinical, and clinical decisionmaking, 48–49

and critical care research, 46–49 to electronic medical record, 41–42 electronic to, 39–40 and clinical research, 42–46

defined, 39

and the electronic medical record,

electronic, 38-39

to clinical databases, 39-40

types, 37

Database management system (DBMS), defined, 39

DBMS. See Database management system. Death, near. See Near-death experience.

Defensive withdrawal, defined, 661

Delirium, causes, 676, 678 cerebral function monitoring for, 756–759 defined, 661, 676

drugs for, 709-731

in ICU, 676-678, 677, 734-735

life-threatening episodes of, 750–753 treatment, 753–755

neurobiology of, 733–734 symptoms, 735–736

treatment, cholecystokinin antagonists, 760-761

drugs in, 733–765. See also Organ system failure. serotonin in, 759–760

Delirium tremens, alcohol withdrawal and, 678, 751

treatment, 753

Depression, in ICU, 677

Dexmedetomidine, for hyperadrenergic hyperfunction, 743

Diabetes mellitus, chronic, conditions associated with, 541–543

coagulation affected by, 542-543

hyperglycemia and, 539–540 hypoglycemia and, 540–541

incidence, 538

infections associated with, 539-540

motor vehicle accidents in patients with, 538

neuropathies associated with, 541-542 in trauma patients, 571

management, 538–543

Dialysis, peritoneal, in traumatized renal failure patients, 530–532

Discomfort, control of, in near-death

experience, 653-654 in near-death experience, 653-654 Discrimination, defined, 56 Distraction, for patient distress during mechanical ventilation, 702 Distress, defined, 661 Distribution half-life, of drugs, 711 Doppler ultrasonography, color, abdominal, 307-309 Doxacurium, description and uses of, 790 Droperidol, pharmacology of, 719-720 Drug(s), for brain failure and delirium, 733-765 inotropic, in trauma patients with pre-existing cardiac disease, 491 overdose of, coma following, 221 in traumatized renal failure patients, 529 Dulin-Williams protocol, description of, 817-819 documentation record, 829 forms used in, 827-829 for monitoring paralyzed patients in respiratory failure, 816-825 nursing checklist, 828 pain and sedation in, 824 physician orders, 827 prolonged immobilization in, 824-825 protocol preparation, 819-824 Dyspnea, defined, 661, 852 in long-term ventilated patients, 852-856, 858-860 Eating, long-term ventilation effects on, 862 ECG. See Electrocardiogram. Echocardiography, in pericardial disease, transesophageal, in trauma patients with

pre-existing peripheral arterial disease, 575 Eclampsia, in pregnant trauma patients, 630-631 Ectopic beats, in trauma patients, treatment, 498 Electrical-mechanical dissociation, in trauma patients, treatment, 498 Electrocardiogram (ECG), in trauma patients with pre-existing cardiac disease, 480 Electrodiagnosis, in evaluation of muscle weakness following neuromuscular blockade, 802-804 Electrolyte(s), abnormalities of, in traumatized renal failure patients, 527-528

Electromyography, needle, in evaluation of muscle weakness following neuromuscular blockade, 803-804 single-fiber (SFEMG), in evaluation of muscle weakness following neuromuscular blockade, 803 Embolism, pulmonary, CT for, 272 screening for, 351 Embolization, treatment of, in the ICU, 446-450 Embolotherapy, percutaneous transcatheter, 446-450 Encephalopathy, hepatic, 563 End-stage renal disease. See also Kidney disease. incidence, 523 Endocrine disease, pre-existing, in trauma patients, management, 537-554. See also specific disease. Endocrine system, pregnancy-induced changes in, 625 Endotracheal tubes, placement and misplacement of, 417-419 Enteric feeding tubes, inadvertent misplacement of, 249 Epilepsy, traumatic injury in patients with, Epinephrine, in trauma patients with preexisting cardiac disease, 491 Esmolol, for life-threatening delirium, 754 for sinus tachycardia, 494 Ethanol, abuse of, in trauma patients, 596-598 Ethic(s), in use of predictive scoring systems, 229-241. See also Scoring system(s), predictive, ethical appropriateness of. Etomidate, pharmacology of, 718 Extracorporeal membrane oxygenator (ECMO), described, 249-250

examination of, 283 Extrathoracic subclavian artery, injury to, in trauma patients with pre-existing peripheral arterial disease, treatment, 583-584 Extremities, lower, venous ultrasonography of, 341-353

Extraluminal air, plain film radiographic

upper, venous ultrasonography of, 353-354

Feeding tubes, enteral, placement and misplacement of, 420-421 enteric, inadvertent misplacement of, 249 nasogastric, placement and misplacement of, 420-421

Fentanyl, for hyperadrenergic hyperfunction, 741-742 in organ system failure, 745-746 transdermal, for pain in mechanically ventilated patients, 768-769 Fluid collections, extrapancreatic, CT of,

infected, percutaneous management of, 445-446

Fluid resuscitation, in trauma patients with pre-existing peripheral arterial disease,

Flumazenil, uses of, 748-749

Gadopentetate dimeglumine, in MRI, 403 Gallbladder, ultrasonography of, 297-303 Gallstone(s), ultrasonographic detection of,

Gas exchange, abnormalities of, in trauma patients with chronic pulmonary disease, 515-516

trauma effects on, 512

Gastrointestinal bleeding, radionuclide evaluation of, 375-378

Gastrointestinal system, narcotic effects on,

pregnancy-induced changes in, 625

Genitourinary tract, calcifications in, plain film radiographic examination of, 293 Goiter, trauma patients with, 549

Graft infection(s), in trauma patients with pre-existing peripheral arterial disease, treatment, 581-582

Graft thromboses, in trauma patients with pre-existing peripheral arterial disease,

Great vessels, renal, ultrasonographic detection of, 312-313

Half-life, of drugs, 711 Haloperidol, pharmacology of, 719-720 Head injury, vegetative state following, outcome from, 224 Head trauma, MRI of, 410-412

Health Policy Corporation of Iowa (HPCI) plan, 147-148

Heart block, complete, in trauma patients, treatment, 498

first-degree, in trauma patients, treatment, 497 second-degree, in trauma patients, treat-

ment, 497-498 Heart disease, ischemic, causes, 474-475 clinical syndromes in, 475-476 pre-existing, in trauma patients, 463

treatment, 485-487 terminology associated with, 474 pre-existing, in trauma patients, 461-506 angina, 466

arrhythmias, 469-470

cardiac risk factors associated with, 465-474

ECG in, 480

evaluation, 474-480

heart failure, 466-468

hemodynamic monitoring for, 480-483 hospital stay affected by, 465

hypertension, 470 incidence, 461-464

mitral insufficiency, 469

mortality rates associated with, 464-465

myocardial infarction, 465-466

significance of, 464-474

treatment, 483-499. See also specific disorder, e.g., Arrhythmias(s). surgical problems, 483-484

valvular heart disease, 468-469

valvular, pre-existing, in trauma patients, 468-469

treatment, 492-493

Heart failure, congestive, defined, 477 diagnosis, 480 pathophysiology, 477-479

in trauma patients, evaluation, 477-480

in trauma patients, 466-468 with pre-existing cardiac disease, treatment, 489-490

Heart rate, control of, in trauma patients with pre-existing cardiac disease, 490-491

Heart valves, prosthetic, in trauma patients with pre-existing cardiac disease, 493 Hematologic system, pregnancy-induced

changes in, 624-625 Hematoma(s), hepatic, ultrasonographic detection of, 306-307

mediastinal, chest CT for, 271

Hemodialysis, in traumatized renal failure patients, 530-532

Hemodynamic instability, sedation problems in, 744

Hemodynamic profiles, in trauma patients with pre-existing cardiac disease, 482-483

Hemopericardium, loculated, 406 Hemorrhage, acute, abdominal, CT of, 335-336

renal, CT of, 331

Hepatobiliary inflammatory disease, acute, CT of, 321-323

Hofmann elimination, 787

Holter monitoring, in trauma patients with pre-existing peripheral arterial disease, Hydromorphone, in organ system failure, 745-746

Hyperglycemia, in trauma patients with diabetes mellitus, 539–540

Hyperosmolar nonketotic coma (HNKC), described, 540

Hypertension, portal, in patients with preexisting liver disease, 560–561

pre-existing, in trauma patients, 463 in trauma patients, 470, 571

evaluation, 476-477

with pre-existing peripheral arterial disease, treatment, 577–578 treatment, 499

Hyperthyroidism, described, 548 management, 549

manifestations of, 548 Hypnosis, for patient distress during mechanical ventilation, 103

Hypoalbuminemia, in patients with preexisting liver disease, 561

Hypoglycemia, in trauma patients with diabetes mellitus, 540–541

Hypotension, in trauma patients with preexisting cardiac disease, 483–484

Hypothyroidism, described, 547 management, 547–548 manifestations of, 548

ICU. See Intensive care unit. Imagery, for patient distress during mechanical ventilation, 103

Imaging, chest, in the ICU, 247–265. See also Chest radiography. in the ICU, 247–454. See also specific

type of imaging.

Immune system, sleep deprivation effects

on, 689 Indomethacin, for preterm labor, 631

Indomethacin, for preterm labor, 631
Infection(s), abdominal, scintigraphy for,
387

CNS, MRI of, 413-414

diabetes mellitus and, 539–540

graft, in trauma patients with pre-existing peripheral arterial disease, treatment, 581–582

pelvic, scintigraphy for, 387

pulmonary, gallium scintigraphy in, 385–386

radiographic evaluation of, 257–260 renal, ultrasonographic detection of, 313–314

wound, in trauma patients with pre-existing peripheral arterial disease, treatment, 581 Infectious diseases, scintigraphy for, 383–391. See also Scintigraphy, for infectious diseases.

sleep and, 682

Inferior vena cava filters, in the ICU, 443-445

placement and misplacement of, 434 Inotropic agents, in trauma patients with pre-existing cardiac disease, 491

Instrumentation, predictive, development of, statistical approaches, 19–35. See also Intensive care unit (ICU) outcome, prediction of, instrumentation for.

Intensive care unit (ICU). See also under Ventilation, mechanical.

anxiety in, 673-675

brain failure in, treatment, drugs in, 733–765. See also *Brain failure*; *Delirium*.

care in, cost of, 2

catheter fragment retrieval in, 441–443 central venous catheter insertion in, 438–441

chest CT in, 267–275. See also Computed tomography (CT), chest.

clinical performance in, 173-174

collaborative practice in, 699 communication with intubated patients, 696–698

competent caring and the nurse-patient relationship in, 698-699

cost-effectiveness in, 173–174 current situations regarding, 169–172 decision-making in, 76–78

delirium in, 676-678

treatment, drugs in, 733-765

duration of stay in, 160–161 emotional tone of, 696–700

fairness in, determination of, 137-138 future trends in, 173-175

identification of low-risk monitor patients in, 161–162

imaging in, issue on, 247-454

inferior vena caval filters in, 443–445 intervention in, 437–452

life and near-death in, 651-657

neurobehavioral abnormalities in, terms used during, 661

neuromuscular blockade in, muscle weakness following, 799–813. See also Neuromuscular blockade, muscle weakness following.

neuromuscular blocking agents in, pharmacology of, 779–797. See also Neuromuscular blocking agents, pharmacology of.

patient distress in, terms used during, 661

percutaneous nephrostomy in, 450–451

quality of care in, recommendations for, 168–169

rationing in, 135-143 avoidance of, 140-142

fair policy for, development of, 139-140

respiratory failure in paralyzed patients in, monitoring and preventive care of, 815–829

sedative reversal agents in, 747-749 sleep determination in, 682-685

sleep disturbances in, **681–694**. See also Sleep disturbances, in ICU.

terminology of, standardization of, 660 therapeutic relationships in, 696–700 recommendation for improvement of, 699–700

utilization of, 136-137, 160-162

vascular ultrasound in, **341–363**. See also *Ultrasonography*, vascular.

Intensive care unit (ICU) outcome, from cardiac arrest, 188–191

from CNS injury, 217-228

critical care medicine effects on, 165–168 evaluation, 2, 160

prediction of. See also Scoring system(s), predictive.

APACHE system for, case against using, 117–125, 127–128, 129–134 collection and analysis of the database, 6–7

databases in, 37-51. See also Database(s).

instrumentation for, development of, multivariate logistic regression analysis, 22–24

> objective methods, 21-22 severity of illness and, 19-20 statistical approaches, 19-24

subjective methods, 20–21 uncertainty associated with, 19–20 validation of, accuracy in, 27–33

model calibration in, 25–27 receiver operating characteristic analysis in, 27–33

issue on, 1-241

model of, development of, 7–9 impact and utility of, 12–13 updates of, 13 validation of, 9–11

models for, severity, 93–110. See also Scoring system(s), predictive.

patient population in, 5–6 scoring systems for, 1–18, 73–89, 91–92. See also Scoring system(s), predic-

tive, in ICU outcome. evaluation, 162-165

in reimbursement strategies, 145-155

in trauma patients, multicenter study for, 120-124

validation of, statistical approaches, 24-33

variables in, 4-5 selection of, 6

predictors of, from critical illness, 179–195

research concerning, 13-15

from respiratory failure, 197–215

from shock, 179-180

Intensive care unit (ICU) psychosis, 736 Intensive care unit (ICU) stress delirium, 735–736

Intensive care unit (ICU) syndrome, defined, 688

sleep deprivation and, 686–689 Intercostal nerve blocks, for pain in mechanically ventilated patients, 774–775

Intestinal diseases, acute, CT of, 334–335 Intestine(s), infarction of, CT of, 335 perforation of, CT of, 334–335

Intra-arotic counterpulsation balloon (IACB), placement and misplacement of, 430–431

Intra-arterial blood pressure, in trauma patients with pre-existing cardiac disease, 481

Iowa, reimbursement strategies in, severity scoring systems for, 146–148

Ischemia, acute mesenteric, plain film radiographic examination of, 283, 284 IVOX, described, 250

Jaundice, CT of, 323–324 scintigraphic evaluation of, 373–374

Ketamine, pharmacology of, 717 Ketoacidosis, diabetic, defined, 540 management, 540 pathogenic factors, 540 Ketorolac, in organ system failure, 745–746

Ketorolac tromethamine, for pain in mechanically ventilated patients, 774

Kidney(s), infarction of, CT of, 330 scintigraphy of, 391–399 transplantation of, scintigraphy for,

392–396 trauma to, ultrasonographic detection of, 314–315

ultrasonography of, 309-315

Kidney disease, acute, CT of, 329–332 pre-existing, in trauma patients, **523–536** incidence, 523 Kidney disease (Continued) prevention, 533 Kidney failure, acute, CT of, 330 scintigraphy for, 396-399 ultrasonographic detection of, 310-312 described, 526-527 in trauma patients wtih pre-existing liver

disease, 563-564

traumatized patients with, blood transfusion in, 528-529 drugs for, 529 electrolyte abnormalities in, 527-528

nutrition for, 529-530 resuscitation following, 528

water and solute removal in, 530-532 Kidney function, impaired, in trauma patients with pre-existing peripheral arterial disease, treatment, 578 trauma effects on, 524-525

Kidney transplantation, incidence, 523 trauma in patients with, 532

Labetalol, for life-threatening delirium, 754 Labor, preterm, management, 631 Lactic acidosis, in shock, 181-183 Linear regression models, predictive, 7-9 Liver, abcesses of, ultrasonographic detection of, 304-305 CT of, 321-323

hematomas of, ultrasonographic detection of, 306-307

ultrasonography of, 304-309 Liver disease, pre-existing, defined, 555

in trauma patients, 555-566 ascites, 560-563 cardiovascular abnormalities, 559-560 coagulopathy, 560

hospital stay affected by, 555-556 mortality affected by, 556-558 pathophysiologic considerations,

558-564 Lorazepam, for hyperadrenergic

hyperfunction, 739-740 in organ system failure, 746 for sleep disturbances in ICU, 690-691

Lung(s), expansion of, in trauma patients with chronic pulmonary disease, 516

Lung disease, chest CT for, 268-269 evaluation of, chest radiography for, 256-260

in trauma patients, 571

Lung injury, acute, pathogenesis, 510

Magnesium sulfate, for preterm labor, 631

Magnetic resonance imaging (MRI), 401-416

advantages of, 401 for AIDS, 413-414

for CNS infections, 413-414

contrast agents for, 403 for head trauma, 410-412

limitations of, 401-403

for musculoskeletal disorders, 409-410 of pelvic abscesses, 409

in pregnant patients, 403

for spinal cord compression, 412-413

for stroke, 410-412

uses, 403-414

for vascular problems, 404-409

Marijuana, abuse of, in trauma patients,

Maternal-fetal "unit." 625-626

Mechanical ventilation. See Ventilation, mechanical.

for acute respiratory failure, prognosis following, 203-204

Mediastinum, abnormalities of, chest CT for, 271

Medical record(s), electronic, clinical databases to, 41-42

databases and, predictive, 38-42 Megacolon, toxic, plain film radiographic

examination of, 283 Mental status, during mechanical

ventilation evaluation, 660-665 Metoprolol, for life-threatening delirium,

754 Midazolam, for hyperadrenergic hyperfunction, 739

in organ system failure, 746 for sleep disturbances in ICU, 690-691

β-Mimetics, for preterm labor, 631 Mitral insufficiency, in trauma patients, 469

Mitral valve disease, in trauma patients with pre-existing cardiac disease, treatment, 493

Mivacurium, for respiratory failure, indications, 840

Morphine, in organ system failure, 745-746 Mortality Prediction Model, errors and bias in, 64

Mortality Prediction Model II, 98-99 Motor vehicle accidents, alcohol abuse and, 595, 596

diabetics and, 538

pregnancy and, 623 MRI. See Magnetic resonance imaging.

99m Technetium-HMPAO, for infection detection, 390

Multivariate logistic regression analysis, predictive, 22-24

Musci therapy, for patient distress during mechanical ventilation, 702-703

Muscle biopsy, in evaluation of muscle weakness following neuromuscular blockade, 804

Muscle relaxants. See also Neuromuscular blocking agents.

administration of, strategies of, 792–797 description and uses of, 785–787 nondepolarizing, description and uses of,

787–790

Musculoskeletal disorders, MRI of, 409–410 Myocardial depression, in trauma patients with pre-existing cardiac disease, treatment, 491

Myocardial infarction, diagnosis, 476 pathophysiology, 475–476 prior, in trauma patients, 465–466

Naloxone, uses of, 748 Narcotic(s), intravenous, abuse of, in trauma patients, 598–600

Nasogastric feeding tubes, placement and misplacement of, 420–421

Near-death experience, discomfort in, 653-654

disease-specific experiences related to, 654-656

in intensive care unit, **651–657** loss of control in, 652–653 pain in, 653–654 patient–physician relationship in, 657 surprising thoughts in, 656

Nerve biopsy, in evaluation of muscle weakness following neuromuscular blockade, 804

Nerve conduction studies, in evaluation of muscle weakness following neuromuscular blockade, 802–803

Neurobehavioral abnormalities, in ICU, terms used during, 661

Neuroleptic agents, pharmacology of, 719–720

Neurologic disorders, traumatic injury in patients with, 635-641 impact of, 638 management, 639-640

risk of, 635–638 Neuromuscular blockade, muscle weakness following, **799–813** assessment, electrophysiology in,

802–804 nerve and muscle biopsy, 804 causes, 805–810

clinical approach to, 800-802 differential diagnosis, 801

persistent, due to drug effect, 805 and reversal of blockade, 781-783

Neuromuscular blocking agents (NMBAs). See also *Muscle relaxants*. administration of, strategies of, 792–797

depolarizing, 781

nondepolarizing, 781 paralysis due to, respiratory failure in patients with, monitoring and preventive care of, 815-829

pharmacodynamics of, 791–792 pharmacokinetics of, 791–792

pharmacology of, 779-797

historical perspective, 779–780 for respiratory failure, 831–843 guidelines for, 839–841

indications for, 832–834 types, 834–839

Neuromuscular junction, anatomy of, 782 and neuromuscular transmission, 780–781

Neuromuscular synapse, 780 Neuromuscular transmission, neuromuscular junction and, 780–781 Neuropathy(ies), diabetes mellitus and,

541-542

Nifedipine, for supraventricular tachycardia, 495–496

Nonintensive care unit outcomes, evaluation, 153–154

Nuclear image(s), production of, principles of, 366

Nuclear medicine, in acutely ill patients, 365–381, 383–400

Nursing assignment(s), in ICU, future trends, 174–175

Nutritional support, in trauma patients with pre-existing peripheral arterial disease, 573

in trauma patients with pre-existing liver disease, 563

in traumatized renal failure patients, 529-530

Obesity, defined, 613
health-related aspects of, 613
in trauma patients, 613–622
airway securing, 614
anatomic considerations, 614–615
cardiovascular considerations, 616–617
conditions associated with, 618–619
indwelling vascular catheter complications, 614–615
intrahospital transport, 615
mechanism of injuries, 613–614
outcome assessment and prediction,

619 respiratory considerations, 615–616 venous thromboembolism in, 617–618 Obesity (Continued)

uncomplicated, respiratory manifestations of, 616

Objective probability, defined, 54

Obstructive lung disease(s), postoperative complications related to, 512–513

Obtundation, defined, 661

Opioid(s), cardiovascular effects, 724

clinical indications, 725-726

dependence on, 726

epidural, continuous infusion of, for pain in mechanically ventilated patients, 773

for pain in mechanically ventilated patients, 771–773

gastrointestinal effects, 724

intravenous, pharmacology of, **709–731** for pain in mechanically ventilated pa-

tients, 770, 771-772

parenteral, for pain in mechanically ventilated patients, 768-769

pharmacokinetics of, 721-723

pharmacology of, 720-726

receptor physiology of, 720-721 respiratory effects, 723

side effects, 723

Organ system failure, analgesics in, 745-746

sedatives in, 746-747

problems related to, 744

Osteomyelitis, detection of, scintigraphy for, 387–390

Pacemaker(s), cardiac, placement and misplacement of, 432–434

Pain, defined, 661

in mechanically ventilated patients, 767–778. See also Ventilation, mechanical, pain associated with, management.

in near-death experience, 653-654 in trauma patients with pre-existing pe-

in trauma patients with pre-existing peripheral arterial disease, treatment, 578

Pancreas, calcifications in, plain film radiographic examination of, 294 Pancreatic inflammatory disease, acute, CT

of, 324–328 Pancreatitis, hemorrhagic, CT of, 327

necrotizing, CT of, 327 Pancuronium, description of, 790 for respiratory failure, 834, 835, 837

uses of, 790
Patient-physician relationship, in neardeath experience, 657

Patient-ventilator asynchrony, defined, 661

Patient-controlled analgesia (PCA), for pain in mechanically ventilated patients, 769

Patient distress, in ICU, terms used during, 661

Patient tolerance, of mechanical ventilation, 659–671. See also *Ventilation*, mechanical.

Pelvis, abscess of, MRI of, 409

infection of, scintigraphy for, 387 Percutaneous intravascular central catheter

(PICC), described, 249
Percutaneous nephrostomy tube

placement, in the ICU, 450–451 Pericardial disease, echocardiography for,

273–274
Peripheral arterial disease (PAD), age as factor in, 567

incidence, 567

pre-existing, in trauma patients, **567–593** antibiotic prophylaxis, **578–579**

aortic reconstruction, 584-588 axillary artery injury, 584

bleeding, treatment, 579–580 brachiocephalic artery injury, 583

cardiac evaluation, 572–573 cardiac function, 574–575

carotid artery injury, 582-583

cerebral circulation, 573

CNS function, 575 diagnosis, 571-573

extrathoracic subclavian artery injury, 583–584

fluid resuscitation for, 576–577 graft infections, treatment, 581–582

graft thrombosis, 580-581 impaired renal function, treatment, 578

laboratory evaluation, 572 monitoring for, 574-576

nonvascular disease, 571

nutritional assessment, 573 pain management, 578

patient history, 572 physical examination, 572

prevalence, 567, 568 pulmonary function, 573

renal insufficiency, 571

significance of, 568-571

treatment, 576-588 vascular complications, treatment, 579-580

ventilatory management, 578

wound infection, treatment, 581 Peripheral vessel patency, monitoring of, in trauma patients with pre-existing peripheral arterial disease, 575–576

Peritoneal dialysis, in traumatized renal failure patients, 530–532

Peritonitis, CT of, 334

Pharmacologic paralysis, of mechanically ventilated patients, 667–668

Phencyclidine, abuse of, in trauma patients, 604-605

Phleboliths, plain film radiographic examination of, 293

Phlegmon(s), CT of, 325

Pituitary—adrenal axis, pre-existing dysfunction of, in trauma patients, 543–545

Pleural disease, chest CT for, 269–270 Pleural drainage tubes, placement and misplacement of, 421–423

Pneumatosis intestinalis, plain film radiographic examination of, 285–287, 290

Pneumocystis carinii pneumonia, respiratory failure associated with, outcome following, 211

Pneumonia, CMV, 257

Pneumocystis carinii, respiratory failure associated with, outcome following, 211

Pneumoperitoneum, plain film radiographic examination of, 283–286

Pneumoretroperitoneum, plain film radiographic examination of, 284–285, 287

Pneumothorax, in ICU patients, diagnosis, 270

Polyneuropathy, critical illness, 806–808 Polysomnography, in sleep measurement, 684, 685

Portal vein, gas in, plain film radiographic examination of, 288, 291

Post-traumatic stress disorder (PTSD), in

Pre-eclampsia, in pregnant trauma patients, 630–631

Pregnancy, maternal-fetal "unit" in, 625–626

trauma in, 623-634

MRI during, 403 physiologic changes of, 624–625 preterm labor during, management, 631

abruptio placentae, 630 amniotic fluid embolus, 630 carbon monoxide poisoning, 631 causes, 623 eclampsia, 630–631 epidemiology, 623–624 fetal assessment, 629–630 maternal assessment, 626–629 outcome, 632 perimortem cesarean section, 631

pre-eclampsia, 630–631 prognosis, 632

uterine rupture, 630 Probability estimate, defined, 54 Procainamide, for atrial fibrillation/flutter, 495–496

Propofol, for life-threatening delirium, 753–754

in organ system failure, 746-747 pharmacology of, 718-719

Propranolol, for atrial fibrillation/flutter, 495

Pseudoaneurysm(s), CT of, 327 Pseudocyst(s), CT of, 325–326

Psychiatric disorders, traumatic injury in patients with, 635-641

impact of, 638 management, 639-640 risk of, 635-638

Psychoactive substance use disorders (PSUDs), diagnosis, 606–607 pitfalls in, 608

prevalence, 607-608

Psychoactive substance withdrawal syndrome, in ICU, 677

Psychological problems, in long-term ventilated patients, 850–852, 857 Psychosis, brief reactive, in ICU, 677

ICU, 736 Pulmonary arterial pressure, evaluation of,

chest radiography for, 253
Pulmonary artery pressure, in trauma
patients with pre-existing cardiac

disease, 481–482 Pulmonary blood flow volume, evaluation of, chest radiography for, 251–253

Pulmonary capillary wedge pressure (PCWP), evaluation of, chest radiography for, 253–255

Pulmonary disease, chest CT for, 268–269 chronic, in trauma patients, **507–522** approach to, 507–510

arterial blood gases, 509-510 chest roentgenography, 510 management, 514-518

obstructive lung disease, 512–513 patient history, 507–508

physical examination, 508 pulmonary function testing, 508, 509 smoking effects on, 512

evaluation of, chest radiography for, 256-260

Pulmonary emboli, CT for, 272 screening for, 351

Pulmonary function testing, for trauma patients with chronic pulmonary disease, 508, 509

Pulmonary infection, gallium scintigraphy in, 385–386

Pyelonephritis, acute, CT of, 329 emphysematous, plain film radiographic

examination of, 289, 292

Quinidine, for atrial fibrillation/flutter, 495 for supraventricular tachycardia, 495–496

Radiography, plain film, of the abdomen, 277–296

Radionuclide scanning, brain, in brain death confirmation, 379

Radiopharmaceuticals, principles of, 365–366

Rationing, in the ICU, 135-143 avoidance of, 140-142

Rehabilitation, of patients on long-term mechanical ventilation, 862–863

Reimbursement, strategies for, ICU severity scoring systems in, 145–155

choosing the quality monitors in, 150–151

in Cleveland, 148-150 in Iowa, 146-148

Relaxation, for patient distress during mechanical ventilation, 701–702

Reliability, defined, 38

Renal disease. See Kidney disease.

Renal insufficiency, in trauma patients, 571 Repetitive nerve stimulation (RNS), in evaluation of muscle weakness following neuromuscular blockade,

803
Reproductive system, pregnancy-induced

changes in, 624
Resource(s), rationing of, in the ICU,
135–143

Respiratory failure, acute, defined, 198–199 mechanical ventilation for, prognosis following, 203–204

outcome following, 199–203

associated with hematologic malignancy, outcome following, 210–211 associated with *Pneumocystis carinii*

associated with *Pneumocystis carini*pneumonia, outcome following, 211
hypoxic/hypercapnic, 199–205
long-term follow-up of patients with,

204–205

ICU outcome from, 197–215 neuromuscular blocking agents for, 831–843. See also Neuromuscular blocking agents (NMBAs), for respira-

tory failure.
paralyzed patient in, monitoring and
preventive care of, 815-829

secondary to status asthmaticus, outcome following, 211–212

in trauma patients with chronic pulmonary disease, 516-518

Respiratory system, benzodiazepine effects on, 713–714 narcotic effects on, 723 in obese trauma patients, 615–616 pregnancy-induced changes in, 625 sleep deprivation effects on, 689 trauma effects on, 510–512

Restraints, for patients on mechanical ventilation, 705

Resuscitation, of traumatized renal failure patients, 528

Ritodrine, for preterm labor, 631 Roentgenography, chest, for trauma patients with chronic pulmonary disease, 510

SAPS, 60

SAPS II system, 98

Scintigraphy, for acute stroke, 379–380 brain, 378–380

in evaluation of gastrointestinal bleeding, 377-378

gallium, for pulmonary infection, 385-386

gastrointestinal, 370 hepatobiliary, 370-375

for infectious diseases, 383–391 agent selection, 383–384

imaging techniques, 383-385

pulmonary, 366-369 availability of, 368

methodology in, 366–368 special problems and concerns with, 369

renal, 391–399

for TIAs, 379–380 Scoring system(s), predictive, **1–18**. See also Intensive care unit (ICU) outcome,

prediction of. in allocation of resources, 85–87 available systems, 94–99

basic concepts, 54–55

caution with, 93–110, 111–115 characteristics, 53–54

in clinical investigation, 84–85

for cost-containment purposes, 103 errors and bias in, 53–72

ethical appropriateness of, 229-241 case study, 230-231

clinical judgment in, 231–232 medical futility associated with, 232–233

evaluation, 162-165

future use of, 238-239 in ICU outcome, 73-89, 91-92

for individual patients, 79–82, 101 for internal quality assessment, 100 for interpretation of a quality compari-

son study, 100–101

models of, 53

based on objective probability estimates, 57–68 with continuous variables, 59–68 with noncontinuous variables, 58–59 based on subjective probability estimates, 55–57 ractical aspects of, 100–101

practical aspects of, 100–101 properties of, 3–4

for quaity assessment, 82-84

reasons for, 74-78

reimbursement strategies in, 145–155 requirements for, 118

for shock, 180-181

for societal and economic concerns, 74-76

uses of, 79-87, 103-108 future, 238-239

in individual patients, 236-238

Sedation, adjuncts to, for brain failure and delirium, 743

choice of, in the ICU, 437-438

in mechanically ventilated patients, 665–666, 668 withdrawal from, problems related to,

749–750 during withdrawal of life support tech-

nology, 755–756 Sedative(s), intravenous, pharmacology of,

709-731 in organ system failure, 746-747 Sedative reversal agents, role of, 747-749

Sepsis, radiographic detection of, 390–391 in trauma patients, 558

Serotonin, in brain failure and delirium, 759–760

Severity system(s). See Scoring system(s). Shock, arterial lactate concentrations in, 181–183

clinical considerations with, 185–186 cytokine markers in, 184–185 hemodynamic indices in, 183–184 ICU outcome related to, 179–180 prognostic indices for, 180–188 scoring systems for, 180–181 states of, 179

treatment, outcome effects, 186–188 Simplified Acute Physiology Score (SAPS),

Simplified Acute Physiology Score (SAPS) II system, 98

Sinus bradycardia, in trauma patients, treatment, 497

Sinus tachycardia, in trauma patients, treatment, 494

Sleep, in ICU, measurement of, 682–685 and infectious disease, 682 nonrapid eye movement (NREM), 681–682

normal, 681-682

rapid eye movement (REM), 681–682 Sleep disturbances, and the ICU syndrome, 686–689

in mechanically ventilated patients, 665, 681-694

causes, 686-688

effects on respiratory and immune function, 689

measurement of, 682-685 treatment, drugs, 690-691

nonpharmacologic, 689–690

Small bowel, obstruction of, plain film radiographic examination of, 280 Smoking, postoperative pulmonary

complications related to, 512

Sodium retention, in patients with preexisting liver disease, 561 Speech, long-term ventilation effects on,

856, 860–862
Spinal cord, compression of, MRI of,

Spleen, abscesses of, CT of, 328–329

Spleen, abscesses of, CT of, 328–329 diseases of, CT of, 328–329 infarcts of, CT of, 328

Statistics, predictive, 19-35

Status asthmaticus, respiratory failure associated with, outcome following, 211–212

Stress response, defined, 661

Stroke, acute, radionuclide scanning in, 379–380

coma following, prognosis, 219–221 MRI of, 410–412

Stupor, defined, 661

Subjective probability, defined, 54

Substance abuse, in brain-injured trauma patients, 636 defined, 595

disorders related to, in trauma patients, 595-611. See also specific drug.

Succinylcholine, description of, 785 for respiratory failure, indications, 840 uses of, 785

Sundown syndrome, 735

Suriclone, in brain failure and delirium, 760–761

Swallowing, long-term ventilation effects on, 856, 862

T–1/2α, of drugs, 711
Tachyarrhythmia(s), in trauma patients, treatment, 494–495
Tachycardia, sinus, in trauma patients.

Tachycardia, sinus, in trauma patients, treatment, 494

supraventricular, in trauma patients, treatment, 495–496 Tachycardia (Continued)

ventricular, in trauma patients, treatment, 496

Temazepam, for sleep disturbances in ICU, 690–691

TENS. See Transcutaneous electrical nerve stimulation.

Terbutaline, for preterm labor, 631

Thrombosis, deep venous, vascular ultrasonography for, 341–353

graft, in trauma patients with pre-existing peripheral arterial disease, 580-581

renal vein, CT of, 331

venous, in obese trauma patients, 617-618

Thyroid disease, in trauma patients, 545–549

Thyroid function, in the critically ill, 545–547

TIAs. See Transient ischemic attacks.

Torsades de pointes, in trauma patients, treatment, 496–497

Tracheostomy tubes, placement and misplacement of, 419

Transcutaneous electrical nerve stimulation (TENS), for pain in mechanically ventilated patients, 769–770

Transfusion, blood. See *Blood transfusion*. Transient ischemic attacks (TIAs), radionuclide scanning in, 379–380

Transplantation, kidney, incidence, 523 trauma in patients with, 532

renal, scintigraphy for, 392-396

Trauma, cardiac disease effects on, 461–641. See also Heart disease, preexisting, in trauma patients.

chest, CT for, 271

chronic pulmonary disease effects on, 507-522. See also Pulmonary disease, chronic, in trauma patients.

evaluation, 525–526

gas exchange effects of, 512 head, MRI of, 410-412

kidney, ultrasonographic detection of, 314–315

in obese patients, 613-622. See also Obesity, in trauma patients.

in patients with a thyroid goiter, 549 in patients with cerebrovascular disease, 569–570

in patients with coronary artery disease, 569

in patients with diabetes mellitus, 571 management, 538–543

in patients with hypertension, 571

in patients with neurologic disorders, 635-641 in patients with pre-existing dysfunction of the pituitary-adrenal axis, 543-545

in patients with pre-existing endocrine disease, 537–554. See also specific disease.

in patients with pre-existing liver disease, 555-566. See also Liver disease, pre-existing, in trauma patients.

in patients with pre-existing peripheral arterial disease, 567-593. See also Peripheral arterial disease (PAD), pre-existing, in trauma patients.

in patients with psychiatric disorders, 635-641

in patients with substance abuse disorders, 595-611. See also specific drug, in patients with thyroid disease, 545-549 pre-existing disease effects on, issue on.

461-641

in pregnancy, **623–634**. See also *Pregnancy*, *trauma in*.

renal disease effects on, **523–536**. See also Kidney disease, pre-existing, in trauma patients.

renal function affected by, 524–525 evaluation, 525–526

respiratory system affected by, 510-512 ventilation effects of, 510-511

TRISS methodology, in prediction of ICU outcome, 119–120

Trizolam, for sleep disturbances in ICU, 690–691

Tumor(s), hematologic, respiratory failure associated with, outcome following, 210-211

Ultrasonography, abdominal, **297–319** for acute calculus cholecystitis, 298–301

> for aspiration, 315–316 biliary tree, 297–303

for choledocholithiasis, 302–303 gallbladder, 297–303

for gallstones, 298

kidneys, 309-315 liver, 304-309

of abdominal aortic aneurysm, 354–357 color Doppler, abdominal, 307–309

duplex, abdominal, 307-309

vascular, **341–363** of arterial catheterization complications, 357–359

of calf veins, 351-353

lower extremity, 341-353 ancillary findings, 351

false-negative studies, 349-350

false-positive studies, 350 technique, 347-349 upper extremity, 353-354 Urinary system, pregnancy-induced changes in, 625 Urinoma, CT of, 331-332 Urosepsis, ultrasonographic detection of, 313-314 Uterine rupture, in pregnant trauma patients, 630

Valve(s), heart, prosthetic, in trauma patients with pre-existing cardiac disease, 493 Vascular disorders, calcifications in, plain film radiographic examination of, 294 MRI of, 404-409 Vecuronium, description of, 790 for respiratory failure, 834-837 indications, 840

uses of, 790 Vegetative state, following head injury, outcome from, 224

Venous thrombosis, in obese trauma patients, 617-618

Ventilation, mechanical. See also under Intensive care unit (ICU). anxiety due to, 673-675

delirium due to, 676-678

long-term, 845-866

categories of patients requiring, 846 cost of, 846 dyspnea related to, 852-856, 858-860 improving tolerance of, 856-863 locations of patients receiving, 846 number of patients receiving, 846 psychological problems related to, 850-852, 857

rehabilitation of patients on, 862-863 tolerance with, problems affecting, 850-856

ventilator used for, 847-849 pain associated with, management, 767-778

epidural techniques, 770-773 local anesthetic techniques, 774-775 peripheral analgesic techniques, 774-775

spinal cord-level techniques, 769-773 subarachnoid techniques, 770

systemic approach, 768-769 TENS, 769-770 patient distress during,

communication with patient, 696-698

treatment, congitive-behavioral strategies, 700-704 nonpharmacologic, 695-708 restraints, 705

visiting policies, 704-705 patient tolerance of, improvement goals, 659-671

evaluation of mental status. 660-665

patient's experiences, 666-667 pharmacologic paralysis, 667-668 sedating drug knowledge. 665-666, 668

sleep deprivation evaluation, 665 standardize terminology, 660

issue on, 651-866

in trauma patients with chronic pulmonary disease, 516-518 weaning from, anxiety due to, 678-679

trauma effects on, 510-511

in trauma patients with pre-existing peripheral arterial disease, treatment,

Ventilator(s), mechanical, for long-term ventilation, 847-849

Visiting policies, for patients on mechanical ventilation, 704-705

Visual isolation, in near-death experience,

Volvulus, plain film radiographic examination of, 282

Water retention, in patients with preexisting liver disease, 561 Withdrawal, defensive, defined, 661 Wolff-Parkinson-White syndrome, in trauma patients, treatment, 495-496 Wound infection(s), in trauma patients

with pre-existing peripheral arterial disease, treatment, 581

Zolpidem, for sleep disturbances in ICU,

Zopiclone, in brain failure and delirium, 760